



# Alternating Permutations

Richard P. Stanley

M.I.T.

# Basic definitions

A sequence  $a_1, a_2, \dots, a_k$  of distinct integers is **alternating** if

$$a_1 > a_2 < a_3 > a_4 < \dots ,$$

and **reverse alternating** if

$$a_1 < a_2 > a_3 < a_4 > \dots .$$

# Euler numbers

$\mathfrak{S}_n$  : symmetric group of all permutations of  
 $1, 2, \dots, n$

# Euler numbers

$\mathfrak{S}_n$  : symmetric group of all permutations of  
 $1, 2, \dots, n$

**Euler number:**

$$\begin{aligned} E_n &= \#\{w \in \mathfrak{S}_n : w \text{ is alternating}\} \\ &= \#\{w \in \mathfrak{S}_n : w \text{ is reverse alternating}\} \end{aligned}$$

(via  $a_1 \cdots a_n \mapsto n + 1 - a_1, \dots, n + 1 - a_n$ )

# Euler numbers

$\mathfrak{S}_n$  : symmetric group of all permutations of  
 $1, 2, \dots, n$

## Euler number:

$$\begin{aligned} E_n &= \#\{w \in \mathfrak{S}_n : w \text{ is alternating}\} \\ &= \#\{w \in \mathfrak{S}_n : w \text{ is reverse alternating}\} \end{aligned}$$

(via  $a_1 \cdots a_n \mapsto n + 1 - a_1, \dots, n + 1 - a_n$ )

E.g.,  $E_4 = 5$  : 2143, 3142, 3241, 4132, 4231

# André's theorem

**Theorem** (Désiré André, 1879)

$$y := \sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

$$\begin{aligned} &= 1 + 1x + 1 \frac{x^2}{2!} + 2 \frac{x^3}{3!} + 5 \frac{x^4}{4!} + 16 \frac{x^5}{5!} \\ &\quad + 61 \frac{x^6}{6!} + 272 \frac{x^7}{7!} + \dots \end{aligned}$$

# André's theorem

**Theorem** (Désiré André, 1879)

$$y := \sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

$$= 1 + 1x + 1 \frac{x^2}{2!} + 2 \frac{x^3}{3!} + 5 \frac{x^4}{4!} + 16 \frac{x^5}{5!} \\ + 61 \frac{x^6}{6!} + 272 \frac{x^7}{7!} + \dots$$

$E_{2n}$  is a **secant number**.

$E_{2n+1}$  is a **tangent number**.

# Proof of André's theorem

$$y := \sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

# Proof of André's theorem

$$y := \sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

Choose  $S \subseteq \{1, 2, \dots, n\}$ , say  $\#S = k$ .

# Proof of André's theorem

$$y := \sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

Choose  $S \subseteq \{1, 2, \dots, n\}$ , say  $\#S = k$ .

Choose a reverse alternating permutation  $u = a_1 a_2 \cdots a_k$  of  $S$ .

# Proof of André's theorem

$$y := \sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

Choose  $S \subseteq \{1, 2, \dots, n\}$ , say  $\#S = k$ .

Choose a reverse alternating permutation  $u = a_1 a_2 \cdots a_k$  of  $S$ .

Choose a reverse alternating permutation  $v = b_1 b_2 \cdots b_{n-k}$  of  $[n] - S$ .

# Proof of André's theorem

$$y := \sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

Choose  $S \subseteq \{1, 2, \dots, n\}$ , say  $\#S = k$ .

Choose a reverse alternating permutation  $u = a_1 a_2 \cdots a_k$  of  $S$ .

Choose a reverse alternating permutation  $v = b_1 b_2 \cdots b_{n-k}$  of  $[n] - S$ .

Let  $w = a_k \cdots a_2 a_1, n + 1, b_1 b_2 \cdots b_{n-k}$ .

# Proof (continued)

$$w = a_k \cdots a_2 a_1, n + 1, b_1 b_2 \cdots b_{n-k}$$

Given  $k$ , there are:

- $\binom{n}{k}$  choices for  $\{a_1, a_2, \dots, a_k\}$
- $E_k$  choices for  $a_1 a_2 \cdots a_k$
- $E_{n-k}$  choices for  $b_1 b_2 \cdots b_{n-k}$ .

# Proof (continued)

$$w = a_k \cdots a_2 a_1, n + 1, b_1 b_2 \cdots b_{n-k}$$

Given  $k$ , there are:

- $\binom{n}{k}$  choices for  $\{a_1, a_2, \dots, a_k\}$
- $E_k$  choices for  $a_1 a_2 \cdots a_k$
- $E_{n-k}$  choices for  $b_1 b_2 \cdots b_{n-k}$ .

We obtain each alternating and reverse alternating  $w \in \mathfrak{S}_{n+1}$  once each.

# Completion of proof

$$\Rightarrow 2E_{n+1} = \sum_{k=0}^n \binom{n}{k} E_k E_{n-k}, \quad n \geq 1$$

# Completion of proof

$$\Rightarrow 2E_{n+1} = \sum_{k=0}^n \binom{n}{k} E_k E_{n-k}, \quad n \geq 1$$

Multiply by  $x^{n+1}/(n+1)!$  and sum on  $n \geq 0$ :

$$2y' = 1 + y^2, \quad y(0) = 1.$$

# Completion of proof

$$\Rightarrow 2E_{n+1} = \sum_{k=0}^n \binom{n}{k} E_k E_{n-k}, \quad n \geq 1$$

Multiply by  $x^{n+1}/(n+1)!$  and sum on  $n \geq 0$ :

$$2y' = 1 + y^2, \quad y(0) = 1.$$

$$\Rightarrow y = \sec x + \tan x.$$

# A new subject?

$$\sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

# A new subject?

$$\sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

**Define**

$$\tan x = \sum_{n \geq 0} E_{2n+1} \frac{x^{2n+1}}{(2n+1)!}$$

$$\sec x = \sum_{n \geq 0} E_{2n} \frac{x^{2n}}{(2n)!}$$

# A new subject?

$$\sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x$$

**Define**

$$\tan x = \sum_{n \geq 0} E_{2n+1} \frac{x^{2n+1}}{(2n+1)!}$$

$$\sec x = \sum_{n \geq 0} E_{2n} \frac{x^{2n}}{(2n)!}$$

⇒ **combinatorial trigonometry**

# Exercises on combinatorial trig.

$$\sec^2 x = 1 + \tan^2 x$$

# Exercises on combinatorial trig.

$$\sec^2 x = 1 + \tan^2 x$$

$$\tan(x + y) = \frac{\tan x + \tan y}{1 - (\tan x)(\tan y)}$$

# Exercises on combinatorial trig.

$$\sec^2 x = 1 + \tan^2 x$$

$$\tan(x + y) = \frac{\tan x + \tan y}{1 - (\tan x)(\tan y)}$$

EC2, Exercise 5.7

# Boustrophedon

**boustrophedon:**

# Boustrophedon

**boustrophedon**: an ancient method of writing in which the lines are inscribed alternately from right to left and from left to right.

# Boustrophedon

**boustrophedon**: an ancient method of writing in which the lines are inscribed alternately from right to left and from left to right.

From Greek *boustrophēdon* (*βουστροφηδόν*), turning like an ox while plowing: *bous*, ox + *strophē*, a turning (from *strephein*, to turn)

# The boustrophedon array

1  
0 → 1  
1 ← 1 ← 0  
0 → 1 → 2 → 2  
5 ← 5 ← 4 ← 2 ← 0  
0 → 5 → 10 → 14 → 16 → 16  
61 ← 61 ← 56 ← 46 ← 32 ← 16 ← 0.  
...

# The boustrophedon array

1  
0 → 1  
1 ← 1 ← 0  
0 → 1 → 2 → 2  
5 ← 5 ← 4 ← 2 ← 0  
0 → 5 → 10 → **14** → 16 → 16  
61 ← 61 ← 56 ← **46** ← **32** ← 16 ← 0.  
...

# Boustrophedon entries

- last term in row  $n$ :  $E_{n-1}$
- sum of terms in row  $n$ :  $E_n$
- $k$ th term in row  $n$ : number of alternating permutations in  $\mathfrak{S}_n$  with first term  $k$ , the **Entringer number**  $E_{n-1,k-1}$ .

# Boustrophedon entries

- last term in row  $n$ :  $E_{n-1}$
- sum of terms in row  $n$ :  $E_n$
- $k$ th term in row  $n$ : number of alternating permutations in  $\mathfrak{S}_n$  with first term  $k$ , the **Entringer number**  $E_{n-1,k-1}$ .


$$\sum_{m \geq 0} \sum_{n \geq 0} E_{m+n, [m, n]} \frac{x^m}{m!} \frac{y^n}{n!} = \frac{\cos x + \sin x}{\cos(x + y)},$$

$$[m, n] = \begin{cases} m, & m + n \text{ odd} \\ n, & m + n \text{ even.} \end{cases}$$


# Some occurrences of Euler numbers

(1)  $E_{2n-1}$  is the number of complete increasing binary trees on the vertex set  $[2n + 1] = \{1, 2, \dots, 2n + 1\}$ .

# Five vertices



# Five vertices



Slightly more complicated for  $E_{2n}$

# Proof for $2n + 1$

$b_1 b_2 \cdots b_m$  : sequence of distinct integers


$$b_i = \min\{b_1, \dots, b_m\}$$

# Proof for $2n + 1$

$b_1 b_2 \cdots b_m$  : sequence of distinct integers


$$b_i = \min\{b_1, \dots, b_m\}$$

Define recursively a binary tree  $T(b_1, \dots, b_m)$  by




# Completion of proof

**Example.** 439172856



# Completion of proof

**Example.** 439172856



Let  $w \in \mathfrak{S}_{2n+1}$ . Then  $T(w)$  is complete if and only if  $w$  is alternating, and the map  $w \mapsto T(w)$  gives the desired bijection.

# Orbits of mergings

(2) Start with  $n$  one-element sets  $\{1\}, \dots, \{n\}$ .

# Orbits of mergings

**(2)** Start with  $n$  one-element sets  $\{1\}, \dots, \{n\}$ .

Merge together two at a time until reaching  $\{1, 2, \dots, n\}$ .

# Orbits of mergings

**(2)** Start with  $n$  one-element sets  $\{1\}, \dots, \{n\}$ .

Merge together two at a time until reaching  $\{1, 2, \dots, n\}$ .

$1-2-3-4-5-6, 12-3-4-5-6, 12-34-5-6$   
 $125-34-6, 125-346, 123456$

# Orbits of mergings

**(2)** Start with  $n$  one-element sets  $\{1\}, \dots, \{n\}$ .

Merge together two at a time until reaching  $\{1, 2, \dots, n\}$ .

$1-2-3-4-5-6, 12-3-4-5-6, 12-34-5-6$   
 $125-34-6, 125-346, 123456$

$\mathfrak{S}_n$  acts on these sequences.

# Orbits of mergings

**(2)** Start with  $n$  one-element sets  $\{1\}, \dots, \{n\}$ .

Merge together two at a time until reaching  $\{1, 2, \dots, n\}$ .

$1-2-3-4-5-6, 12-3-4-5-6, 12-34-5-6$   
 $125-34-6, 125-346, 123456$

$\mathfrak{S}_n$  acts on these sequences.

**Theorem.** *The number of  $\mathfrak{S}_n$ -orbits is  $E_{n-1}$ .*

# Orbits of mergings

**(2)** Start with  $n$  one-element sets  $\{1\}, \dots, \{n\}$ .

Merge together two at a time until reaching  $\{1, 2, \dots, n\}$ .

$1-2-3-4-5-6, 12-3-4-5-6, 12-34-5-6$   
 $125-34-6, 125-346, 123456$

$\mathfrak{S}_n$  acts on these sequences.

**Theorem.** *The number of  $\mathfrak{S}_n$ -orbits is  $E_{n-1}$ .*

**Proof.**

# Orbits of mergings

**(2)** Start with  $n$  one-element sets  $\{1\}, \dots, \{n\}$ .

Merge together two at a time until reaching  $\{1, 2, \dots, n\}$ .

$1-2-3-4-5-6, 12-3-4-5-6, 12-34-5-6$   
 $125-34-6, 125-346, 123456$

$\mathfrak{S}_n$  acts on these sequences.

**Theorem.** *The number of  $\mathfrak{S}_n$ -orbits is  $E_{n-1}$ .*

**Proof.** Exercise.

# Orbit representatives for $n = 5$

12-3-4-5

123-4-5

1234-5

12-3-4-5

123-4-5

123-45

12-3-4-5

12-34-5

125-34

12-3-4-5

12-34-5

12-345

12-3-4-5

12-34-5

1234-5

# Volume of a polytope

(3) Let  $\mathcal{E}_n$  be the convex polytope in  $\mathbb{R}^n$  defined by

$$\begin{aligned}x_i &\geq 0, & 1 \leq i \leq n \\x_i + x_{i+1} &\leq 1, & 1 \leq i \leq n - 1.\end{aligned}$$

# Volume of a polytope

(3) Let  $\mathcal{E}_n$  be the convex polytope in  $\mathbb{R}^n$  defined by

$$\begin{aligned}x_i &\geq 0, & 1 \leq i \leq n \\x_i + x_{i+1} &\leq 1, & 1 \leq i \leq n - 1.\end{aligned}$$

**Theorem.** *The volume of  $\mathcal{E}_n$  is  $E_n/n!$ .*

# Naive proof

$$\text{vol}(\mathcal{E}_n) = \int_{x_1=0}^1 \int_{x_2=0}^{1-x_1} \int_{x_3=0}^{1-x_2} \cdots \int_{x_n=0}^{1-x_{n-1}} dx_1 dx_2 \cdots dx_n$$

# Naive proof

$$\text{vol}(\mathcal{E}_n) = \int_{x_1=0}^1 \int_{x_2=0}^{1-x_1} \int_{x_3=0}^{1-x_2} \cdots \int_{x_n=0}^{1-x_{n-1}} dx_1 dx_2 \cdots dx_n$$

$$f_n(t) := \int_{x_1=0}^t \int_{x_2=0}^{1-x_1} \int_{x_3=0}^{1-x_2} \cdots \int_{x_n=0}^{1-x_{n-1}} dx_1 dx_2 \cdots dx_n$$

# Naive proof

$$\text{vol}(\mathcal{E}_n) = \int_{x_1=0}^1 \int_{x_2=0}^{1-x_1} \int_{x_3=0}^{1-x_2} \cdots \int_{x_n=0}^{1-x_{n-1}} dx_1 dx_2 \cdots dx_n$$

$$f_n(t) := \int_{x_1=0}^t \int_{x_2=0}^{1-x_1} \int_{x_3=0}^{1-x_2} \cdots \int_{x_n=0}^{1-x_{n-1}} dx_1 dx_2 \cdots dx_n$$

$$f'_n(t) = \int_{x_2=0}^{1-t} \int_{x_3=0}^{1-x_2} \cdots \int_{x_n=0}^{1-x_{n-1}} dx_2 dx_3 \cdots dx_n$$

$$= f_{n-1}(1-t).$$

$F(y)$

$$f'_n(t) = f_{n-1}(1-t), \quad f_0(t) = 1, \quad f_n(0) = 0 \quad (n > 0)$$

# $F(y)$

$$f'_n(t) = f_{n-1}(1-t), \quad f_0(t) = 1, \quad f_n(0) = 0 \quad (n > 0)$$

$$\mathbf{F(y)} = \sum_{n \geq 0} f_n(t) y^n$$

$$\Rightarrow \frac{\partial^2}{\partial t^2} F(y) = -y^2 F(y),$$

etc.

# Conclusion of proof

$$F(y) = (\sec y)(\cos(t - 1)y + \sin ty)$$

$$\Rightarrow F(y)|_{t=1} = \sec y + \tan y.$$

# Tridiagonal matrices

An  $n \times n$  matrix  $M = (m_{ij})$  is **tridiagonal** if  $m_{ij} = 0$  whenever  $|i - j| \geq 2$ .

**doubly-stochastic**:  $m_{ij} \geq 0$ , row and column sums equal 1

$\mathcal{T}_n$ : set of  $n \times n$  tridiagonal doubly stochastic matrices

# Polytope structure of $\mathcal{T}_n$

**Easy fact:** the map

$$\begin{aligned}\mathcal{T}_n &\rightarrow \mathbb{R}^{n-1} \\ M &\mapsto (m_{12}, m_{23}, \dots, m_{n-1,n})\end{aligned}$$

is a (linear) bijection from  $\mathcal{T}$  to  $\mathcal{E}_n$ .

# Polytope structure of $\mathcal{T}_n$

**Easy fact:** the map

$$\begin{aligned}\mathcal{T}_n &\rightarrow \mathbb{R}^{n-1} \\ M &\mapsto (m_{12}, m_{23}, \dots, m_{n-1,n})\end{aligned}$$

is a (linear) bijection from  $\mathcal{T}$  to  $\mathcal{E}_n$ .

**Application** (**Diaconis** et al.): random doubly stochastic tridiagonal matrices and random walks on  $\mathcal{T}_n$

# A modification

Let  $\mathcal{F}_n$  be the convex polytope in  $\mathbb{R}^n$  defined by

$$\begin{aligned}x_i &\geq 0, & 1 \leq i \leq n \\x_i + x_{i+1} + x_{i+2} &\leq 1, & 1 \leq i \leq n - 2.\end{aligned}$$

$$V_n = \text{vol}(\mathcal{F}_n)$$

# A modification

Let  $\mathcal{F}_n$  be the convex polytope in  $\mathbb{R}^n$  defined by

$$\begin{aligned}x_i &\geq 0, \quad 1 \leq i \leq n \\x_i + x_{i+1} + x_{i+2} &\leq 1, \quad 1 \leq i \leq n - 2.\end{aligned}$$

$$V_n = \text{vol}(\mathcal{F}_n)$$

$n$	1-3	4	5	6	7	8	9	10
$n!V_n$	1	2	5	14	47	182	786	3774

# A “naive” recurrence

$$V_n = \mathbf{f}_n(1, 1),$$

where

$$f_0(a, b) = 1, \quad f_n(0, b) = 0 \text{ for } n > 0$$

$$\frac{\partial}{\partial a} f_n(a, b) = f_{n-1}(b - a, 1 - a).$$

# $f_n(a, b)$ for $n \leq 3$

$$f_1(a, b) = a$$

$$f_2(a, b) = \frac{1}{2}(2ab - a^2)$$

$$f_3(a, b) = \frac{1}{6}(a^3 - 3a^2 - 3ab^2 + 6ab)$$

# $f_n(a, b)$ for $n \leq 3$

$$f_1(a, b) = a$$

$$f_2(a, b) = \frac{1}{2}(2ab - a^2)$$

$$f_3(a, b) = \frac{1}{6}(a^3 - 3a^2 - 3ab^2 + 6ab)$$

Is there a “nice” generating function for  $f_n(a, b)$  or  $V_n = f_n(1, 1)$ ?

# Distribution of $\text{is}(w)$

$\text{is}(w)$  = length of longest increasing subsequence of  $w \in \mathfrak{S}_n$

# Distribution of $\text{is}(w)$

$\text{is}(w)$  = length of longest increasing subsequence of  $w \in \mathfrak{S}_n$

$$\text{is}(48361572) = 3$$

# Distribution of $\text{is}(w)$

$\text{is}(w)$  = length of longest increasing subsequence of  $w \in \mathfrak{S}_n$

$$\text{is}(48361572) = 3$$

# Distribution of $\text{is}(w)$

$\text{is}(w)$  = length of longest increasing subsequence of  $w \in \mathfrak{S}_n$

$$\text{is}(48361572) = 3$$

**Vershik-Kerov, Logan-Shepp:**

$$\begin{aligned} E(n) &:= \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} \text{is}(w) \\ &\sim 2\sqrt{n} \end{aligned}$$

# Limiting distribution of $\text{is}(w)$

## Baik-Deift-Johansson:

For fixed  $t \in \mathbb{R}$ ,

$$\lim_{n \rightarrow \infty} \text{Prob} \left( \frac{\text{is}_n(w) - 2\sqrt{n}}{n^{1/6}} \leq t \right) = F(t),$$

the **Tracy-Widom distribution**.

# Alternating analogues

- Length of longest alternating subsequence of  $w \in \mathfrak{S}_n$

# Alternating analogues

- Length of longest alternating subsequence of  $w \in \mathfrak{S}_n$
- Length of longest increasing subsequence of an alternating permutation  $w \in \mathfrak{S}_n$ .

# Alternating analogues

- Length of longest alternating subsequence of  $w \in \mathfrak{S}_n$
- Length of longest increasing subsequence of an alternating permutation  $w \in \mathfrak{S}_n$ .

The first is much easier!

# Longest alternating subsequences

$as(w)$  = length of longest alt. subseq. of  $w$

$$w = 56218347 \Rightarrow as(w) = 5$$

# Longest alternating subsequences

$as(w)$  = length of longest alt. subseq. of  $w$

$$w = 56218347 \Rightarrow as(w) = 5$$

$$D(n) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} as(w) \sim ?$$

# Definitions of $a_k(n)$ and $b_k(n)$

$$a_k(n) = \#\{w \in \mathfrak{S}_n : \text{as}(w) = k\}$$

$$b_k(n) = a_1(n) + a_2(n) + \cdots + a_k(n)$$

$$= \#\{w \in \mathfrak{S}_n : \text{as}(w) \leq k\}$$

# The case $n = 3$

$w$	$as(w)$
123	1
132	2
213	3
231	2
312	3
321	2

# The case $n = 3$

$w$	$\text{as}(w)$
123	1
132	2
213	3
231	2
312	3
321	2

$$a_1(3) = 1, a_2(3) = 3, a_3(3) = 2$$

$$b_1(3) = 1, b_2(3) = 4, b_3(3) = 6$$

# The main lemma

**Lemma.**  $\forall w \in \mathfrak{S}_n \exists$  *alternating subsequence of maximal length that contains  $n$ .*

# The main lemma

**Lemma.**  $\forall w \in \mathfrak{S}_n \exists$  *alternating subsequence of maximal length that contains  $n$ .*

**Corollary.**

$$\Rightarrow a_k(n) = \sum_{j=1}^n \binom{n-1}{j-1}$$

$$\sum_{2r+s=k-1} (a_{2r}(j-1) + a_{2r+1}(j-1)) a_s(n-j)$$

# The main generating function

$$B(x, t) = \sum_{k, n \geq 0} b_k(n) t^k \frac{x^n}{n!}$$

**Theorem.**

$$B(x, t) = \frac{2/\rho}{1 - \frac{1-\rho}{t} e^{\rho x}} - \frac{1}{\rho},$$

where  $\rho = \sqrt{1 - t^2}$ .

# Formulas for $b_k(n)$

## Corollary.

$$\Rightarrow b_1(n) = 1$$

$$b_2(n) = n$$

$$b_3(n) = \frac{1}{4}(3^n - 2n + 3)$$

$$b_4(n) = \frac{1}{8}(4^n - (2n - 4)2^n)$$

⋮

# Formulas for $b_k(n)$

## Corollary.

$$\Rightarrow b_1(n) = 1$$

$$b_2(n) = n$$

$$b_3(n) = \frac{1}{4}(3^n - 2n + 3)$$

$$b_4(n) = \frac{1}{8}(4^n - (2n - 4)2^n)$$

⋮

no such formulas for longest **increasing** subsequences

# Mean (expectation) of $\text{as}(w)$

$$D(n) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} \text{as}(w) = \frac{1}{n!} \sum_{k=1}^n k \cdot a_k(n),$$

the **expectation** of  $\text{as}(w)$  for  $w \in \mathfrak{S}_n$

# Mean (expectation) of $as(w)$

$$D(n) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} as(w) = \frac{1}{n!} \sum_{k=1}^n k \cdot a_k(n),$$

the **expectation** of  $as(w)$  for  $w \in \mathfrak{S}_n$

Let

$$\begin{aligned} A(x, t) &= \sum_{k, n \geq 0} a_k(n) t^k \frac{x^n}{n!} = (1-t)B(x, t) \\ &= (1-t) \left( \frac{2/\rho}{1 - \frac{1-\rho}{t} e^{\rho x}} - \frac{1}{\rho} \right). \end{aligned}$$

# Formula for $D(n)$

$$\begin{aligned}\sum_{n \geq 0} D(n)x^n &= \frac{\partial}{\partial t} A(x, 1) \\ &= \frac{6x - 3x^2 + x^3}{6(1-x)^2} \\ &= x + \sum_{n \geq 2} \frac{4n+1}{6} x^n.\end{aligned}$$

# Formula for $D(n)$

$$\begin{aligned}\sum_{n \geq 0} D(n)x^n &= \frac{\partial}{\partial t} A(x, 1) \\ &= \frac{6x - 3x^2 + x^3}{6(1-x)^2} \\ &= x + \sum_{n \geq 2} \frac{4n+1}{6} x^n.\end{aligned}$$

$$\Rightarrow D(n) = \frac{4n+1}{6}, \quad n \geq 2$$

# Formula for $D(n)$

$$\begin{aligned}\sum_{n \geq 0} D(n)x^n &= \frac{\partial}{\partial t} A(x, 1) \\ &= \frac{6x - 3x^2 + x^3}{6(1-x)^2} \\ &= x + \sum_{n \geq 2} \frac{4n+1}{6} x^n.\end{aligned}$$

$$\Rightarrow D(n) = \frac{4n+1}{6}, \quad n \geq 2$$

Compare  $E(n) \sim 2\sqrt{n}$ .

# Variance of $as(w)$

$$V(n) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} \left( as(w) - \frac{4n+1}{6} \right)^2, \quad n \geq 2$$

the **variance** of  $as(w)$  for  $w \in \mathfrak{S}_n$

# Variance of $as(w)$

$$V(n) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} \left( as(w) - \frac{4n+1}{6} \right)^2, \quad n \geq 2$$

the **variance** of  $as(w)$  for  $w \in \mathfrak{S}_n$

**Corollary.**

$$V(n) = \frac{8}{45}n - \frac{13}{180}, \quad n \geq 4$$

# Variance of $as(w)$

$$V(n) = \frac{1}{n!} \sum_{w \in \mathfrak{S}_n} \left( as(w) - \frac{4n+1}{6} \right)^2, \quad n \geq 2$$

the **variance** of  $as(w)$  for  $w \in \mathfrak{S}_n$

**Corollary.**

$$V(n) = \frac{8}{45}n - \frac{13}{180}, \quad n \geq 4$$

similar results for higher moments

# A new distribution?

$$P(t) = \lim_{n \rightarrow \infty} \text{Prob}_{w \in \mathfrak{S}_n} \left( \frac{\text{as}(w) - 2n/3}{\sqrt{n}} \leq t \right)$$

# A new distribution?

$$P(t) = \lim_{n \rightarrow \infty} \text{Prob}_{w \in \mathfrak{S}_n} \left( \frac{\text{as}(w) - 2n/3}{\sqrt{n}} \leq t \right)$$

Stanley distribution?

# Limiting distribution

**Theorem** (Pemantle, Widom, (Wilf)).

$$\lim_{n \rightarrow \infty} \text{Prob}_{w \in \mathfrak{S}_n} \left( \frac{\text{as}(w) - 2n/3}{\sqrt{n}} \leq t \right)$$

$$= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{t\sqrt{45}/4} e^{-s^2} ds$$

(Gaussian distribution)

# Limiting distribution

**Theorem** (Pemantle, Widom, (Wilf)).

$$\lim_{n \rightarrow \infty} \text{Prob}_{w \in \mathfrak{S}_n} \left( \frac{\text{as}(w) - 2n/3}{\sqrt{n}} \leq t \right)$$

$$= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{t\sqrt{45}/4} e^{-s^2} ds$$

(Gaussian distribution)



# Umbral enumeration

**Umbral formula:** involves  $E^k$ , where  $E$  is an indeterminate (the **umbra**). Replace  $E^k$  with the Euler number  $E_k$ . (Technique from 19th century, modernized by **Rota** et al.)

# Umbral enumeration

**Umbral formula:** involves  $E^k$ , where  $E$  is an indeterminate (the **umbra**). Replace  $E^k$  with the Euler number  $E_k$ . (Technique from 19th century, modernized by **Rota** et al.)

**Example.**

$$\begin{aligned}(1 + E^2)^3 &= 1 + 3E^2 + 3E^4 + E^6 \\ &= 1 + 3E_2 + 3E_4 + E_6 \\ &= 1 + 3 \cdot 1 + 3 \cdot 5 + 61 \\ &= 80\end{aligned}$$

# Another example

$$\begin{aligned}(1+t)^E &= 1 + Et + \binom{E}{2}t^2 + \binom{E}{3}t^3 + \dots \\ &= 1 + Et + \frac{1}{2}E(E-1)t^2 + \dots \\ &= 1 + E_1t + \frac{1}{2}(E_2 - E_1)t^2 + \dots \\ &= 1 + t + \frac{1}{2}(1-1)t^2 + \dots \\ &= 1 + t + O(t^3).\end{aligned}$$

# An umbral quiz

Let  $B$  be the Bell number umbra. Then

# An umbral quiz

Let  $B$  be the Bell number umbra. Then

$$(1 + t)^B = ??$$

# An umbral quiz

Let  $B$  be the Bell number umbra. Then

$$(1 + t)^B = e^t$$

# Alt. fixed-point free involutions

**fixed point free involution**  $w \in \mathfrak{S}_{2n}$ : all cycles of length two (number =  $1 \cdot 3 \cdot 5 \cdots (2n - 1)$ )

# Alt. fixed-point free involutions

**fixed point free involution**  $w \in \mathfrak{S}_{2n}$ : all cycles of length two (number =  $1 \cdot 3 \cdot 5 \cdots (2n - 1)$ )

Let  $f(n)$  be the number of **alternating** fixed-point free involutions in  $\mathfrak{S}_{2n}$ .

# Alt. fixed-point free involutions

**fixed point free involution**  $w \in \mathfrak{S}_{2n}$ : all cycles of length two (number =  $1 \cdot 3 \cdot 5 \cdots (2n - 1)$ )

Let  $f(n)$  be the number of **alternating** fixed-point free involutions in  $\mathfrak{S}_{2n}$ .

$$n = 3 : \quad 214365 = (1, 2)(3, 4)(5, 6)$$

$$645231 = (1, 6)(2, 4)(3, 5)$$

$$f(3) = 2$$

# An umbral theorem

**Theorem.**

$$F(x) = \sum_{n \geq 0} f(n)x^n$$

# An umbral theorem

**Theorem.**

$$\begin{aligned} F(x) &= \sum_{n \geq 0} f(n)x^n \\ &= \left( \frac{1+x}{1-x} \right)^{(E^2+1)/4} \end{aligned}$$

# Proof idea

**Proof.** Uses representation theory of the symmetric group  $\mathfrak{S}_n$ .

# Proof idea

**Proof.** Uses representation theory of the symmetric group  $\mathfrak{S}_n$ .

There is a character  $\chi$  of  $\mathfrak{S}_n$  (due to **H. O. Foulkes**) such that for all  $w \in \mathfrak{S}_n$ ,

$$\chi(w) = 0 \text{ or } \pm E_k.$$

# Proof idea

**Proof.** Uses representation theory of the symmetric group  $\mathfrak{S}_n$ .

There is a character  $\chi$  of  $\mathfrak{S}_n$  (due to **H. O. Foulkes**) such that for all  $w \in \mathfrak{S}_n$ ,

$$\chi(w) = 0 \text{ or } \pm E_k.$$

Now use known results on combinatorial properties of characters of  $\mathfrak{S}_n$ .

# Ramanujan's Second Notebook

**Theorem** (Ramanujan, Berndt, implicitly) As  $x \rightarrow 0+$ ,

$$2 \sum_{n \geq 0} \left( \frac{1-x}{1+x} \right)^{n(n+1)} \sim \sum_{k \geq 0} f(k) x^k = F(x),$$

an **analytic** (non-formal) identity.

# A formal identity

**Corollary** (via Ramanujan, Andrews).

$$F(x) = 2 \sum_{n \geq 0} q^n \frac{\prod_{j=1}^n (1 - q^{2j-1})}{\prod_{j=1}^{2n+1} (1 + q^j)},$$

where  $q = \left(\frac{1-x}{1+x}\right)^{2/3}$ , a **formal identity**.

# Simple result, hard proof

**Recall:** number of  $n$ -cycles in  $\mathfrak{S}_n$  is  $(n - 1)!$ .

# Simple result, hard proof

**Recall:** number of  $n$ -cycles in  $\mathfrak{S}_n$  is  $(n - 1)!$ .

**Theorem.** Let  $b(n)$  be the number of **alternating**  $n$ -cycles in  $\mathfrak{S}_n$ . Then if  $n$  is odd,

$$b(n) = \frac{1}{n} \sum_{d|n} \mu(d) (-1)^{(d-1)/2} E_{n/d}.$$

# Special case

**Corollary.** *Let  $p$  be an odd prime. Then*

$$b(p) = \frac{1}{p} \left( E_p - (-1)^{(p-1)/2} \right).$$

# Special case

**Corollary.** *Let  $p$  be an odd prime. Then*

$$b(p) = \frac{1}{p} \left( E_p - (-1)^{(p-1)/2} \right).$$

Combinatorial proof?

# Inc. subsequences of alt. perms.

**Recall:**  $\text{is}(w)$  = length of longest increasing subsequence of  $w \in \mathfrak{S}_n$ . Define

$$C(n) = \frac{1}{E_n} \sum_w \text{is}(w),$$

where  $w$  ranges over all  $E_n$  alternating permutations in  $\mathfrak{S}_n$ .

$\beta$

Crude estimate: what is

$$\beta = \lim_{n \rightarrow \infty} \frac{\log C(n)}{\log n} ?$$

I.e.,  $C(n) = n^{\beta+o(1)}$ .

$\beta$

Crude estimate: what is

$$\beta = \lim_{n \rightarrow \infty} \frac{\log C(n)}{\log n} ?$$

I.e.,  $C(n) = n^{\beta+o(1)}$ .

**J. Shearer**, 2010:  $\beta = \frac{1}{2}$ .

# Limiting distribution?

What is the (suitably scaled) limiting distribution of  $\text{is}(w)$ , where  $w$  ranges over all alternating permutations in  $\mathfrak{S}_n$ ?

# Limiting distribution?

What is the (suitably scaled) limiting distribution of  $\text{is}(w)$ , where  $w$  ranges over all alternating permutations in  $\mathfrak{S}_n$ ?

Is it the Tracy-Widom distribution?

# Limiting distribution?

What is the (suitably scaled) limiting distribution of  $\text{is}(w)$ , where  $w$  ranges over all alternating permutations in  $\mathfrak{S}_n$ ?

Is it the Tracy-Widom distribution?

**Possible tool:**  $\exists$  “umbral analogue” of Gessel’s determinantal formula.

# Limiting distribution?

What is the (suitably scaled) limiting distribution of  $\text{is}(w)$ , where  $w$  ranges over all alternating permutations in  $\mathfrak{S}_n$ ?

Is it the Tracy-Widom distribution?

**Possible tool:**  $\exists$  “umbral analogue” of Gessel’s determinantal formula.

.....

**Darn!**

That's  
the  
end...

